

DIALOG(R) File 351: Derwent WPI

(c) 2006 The Thomson Corporation. All rts. reserv.

0012429563

WPI ACC NO: 2002-374715/ XRAM Acc No: C2002-106052

Hair-treating agent composition, contains fluorescent whitening dye and benzylalcohol, N-methyl-2-pyrrolidone and/or ethylene carbonate derivatives as penetration-promoting agent

Patent Assignee: KAO CORP (KAOS)

Inventor: HAMADA H
1 patents, 1 countries

Patent Family

Patent Application

Number Kind Date Number Kind Date Update JP 2001302473 A 20011031 JP 2000119908 A 20000420 200241 B

Priority Applications (no., kind, date): JP 2000119908 A 20000420

## Patent Details

Number Kind Lan Pg Dwg Filing Notes JP 2001302473 A JA 5 0

## Alerting Abstract JP A

NOVELTY - A hair-treating agent composition comprises a fluorescent whitening dye and at least one compound selected from benzylalcohol (I), N-methyl-2-pyrrolidone (II) and ethylene carbonate (III) derivatives as penetration-promoting solvent.

DESCRIPTION - A hair-treating agent composition comprises a fluorescent whitening dye and a benzylalcohol of formula (I), N-methyl-2-pyrrolidone of formula (II) and/or ethylene carbonate of formula (III) or their derivatives as penetration-promoting solvent.

http://imagesrv.dialog.com/imanager/getimage?ref=I9921ab2063b211da99f600008361346f&f=351&type=PNG

R1= H, methyl or methoxy;

R2= single bond, methylene or ethylene;

R3= H, methyl or ethyl;

a = 0 - 2;

R4= 1-18C, optionally branched alkyl;
and

R5= H, methyl or ethyl.

USE - The composition is optionally available and formed in liquid, gel, cream, emulsion, spray or foam form.

ADVANTAGE - The composition allows the fluorescent whitening dye to penetrate into hair efficiently and prevents bleeding of dyes from dyed hair, resulting in improved dyeing performance and fastness after shampooing.

## Technology Focus

ORGANIC CHEMISTRY - Preferred Components: The fluorescent whitening dye is selected from distilbene, benzooxazole, naphthalimide, benzimidazole, pyrazoline and coumalin derivatives.

Compound (I) is preferably benzyl alcohol, 2-benzyloxyethanol, phenoxyethanol, phenoxy isopropanol and methylphenoxyethanol.

Compound (II) is N-methyl -2-pyrrolidone.

Compound (III) is preferably selected from ethylene and propylene carbonate.